

**MR. ASHOK BOHRA MBBS MS MPhil FRCSEd FRCS (Gen.Surg.)
CONSULTANT GENERAL / LAPAROSCOPIC/ UPPER GI SURGEON**

www.ashokbohra.co.uk

Email: enquiries@ashokbohra.co.uk

UG09lite Laparoscopic Nissen Fundoplication

Expires end of December 2011
Issued December 2010
Copyright © 2010 EIDO Healthcare Ltd

You can get more information about this procedure from **www.aboutmyhealth.org**
Tell us how useful you found this document at **www.patientfeedback.org**



www.rcsed.ac.uk



www.asgbi.org.uk



This document will give you information about a laparoscopic Nissen fundoplication. If you have any questions, you should ask your GP or other relevant health professional.

What is acid reflux?

Acid reflux is a condition where acid from the stomach travels up into the oesophagus (gullet). This happens if the valve between the stomach and the oesophagus does not work effectively. Acid reflux can cause 'heartburn' or acid in the back of the mouth.

What are the benefits of surgery?

Surgery is aimed at curing the symptoms of acid reflux.

Are there any alternatives to surgery?

Drugs that lower the acid content in the stomach are effective at controlling symptoms and healing inflammation.

Surgery is recommended only if the symptoms continue or if you would rather have an operation than take medication for the rest of your life.

What does the operation involve?

The operation is performed under a general anaesthetic and usually takes between one and two hours.

The laparoscopic ('keyhole') technique is usually used. Your surgeon will make several small cuts on your abdomen. They will place surgical instruments, along with a telescope, inside your abdomen and perform the operation (see figure 1).

Your surgeon will stitch the diaphragm to reduce the size of the hole the oesophagus passes through. They will then wrap and stitch the top part of the stomach around the lower oesophagus.

The abdominal cavity is inflated with carbon dioxide gas to help with surgery

A telescope and instruments allow your surgeon to examine the abdomen




Figure 1

The technique for laparoscopic surgery

What complications can happen?

1 General complications

- Pain
- Bleeding
- Unsightly scarring
- Developing a hernia in the scar
- Infection of the surgical site (wound)
- Blood clots

2 Specific complications

- Damage to internal organs
- Developing a hernia near one of the cuts
- Surgical emphysema
- Difficulty swallowing
- Air in the chest cavity
- Making a hole in the oesophagus or stomach
- Tear of the stitches used for the wrap
- Damage to the liver

Long-term problems

- Incomplete control of reflux symptoms
- Weight loss
- Abdominal discomfort
- Diarrhoea

How soon will I recover?

You should be able to go home the next day. You should be able to return to work after three to four weeks but this may vary depending on the extent of surgery and your type of work.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, you should ask a member of the healthcare team or your GP for advice.

You should make a full recovery, with the symptoms of acid reflux gone or much improved.

Summary

Acid reflux can cause heartburn or acid in the mouth. The acid can cause the lining of the oesophagus to become inflamed or scarred. Surgery may be recommended if your symptoms continue while you are on medication.

Acknowledgements

Author: Mr Simon Parsons DM FRCS (Gen. Surg.)
Illustrations: Medical Illustration Copyright © 2010
Nucleus Medical Art. All rights reserved.
www.nucleusinc.com

This document is intended for information purposes only and should not replace advice that your relevant health professional would give you.